Tensor network and a black hole

石原雅文

東北大学原子分子材料科学高等研究機構 (WPI-AIMR) Collaborators: 松枝宏明 仙台高等専門学校 橋爪洋一郎 東京理科大学

> H. Matsueda, M. I. and Y. Hashizume Phys. Rev. D 87, 066002 (2013) arXiv:1208.0206

Introduction

Recently the relation between Tensor Network of wave function in quantum critical phase and discrete Anti de Sitter (AdS) space has been suggested. (B.Swingle '2009)



MERA: (Multiscale Entanglement Renormalization Ansatz):

tensor network describing the wave function in critical phase

Outline

1 Tensor Network (MPS, MERA)

2 Tensor Network and Entanglement Entropy

3 AdS/CFT and MERA

4 MERA at thermal system and AdS black hole



Tensor Network: graphical representation of the wave function of quantum many-body system

Matrix Product States (MPS)

External line: spin indices Internal line: inner product

E.g. Tensor Network in 2-site Heisenberg model

2-site anti-ferromagnetic Heisenberg model.

$$H = J\overrightarrow{S_1} \cdot \overrightarrow{S_2} = \frac{1}{2}J(S_1^+S_2^-) + JS_1^zS_2^z$$

Ground state and its MPS representation

A, B are the following vectors.

$$A^{\uparrow}=(\mathbf{1},\mathbf{0}), \ A^{\downarrow}=(\mathbf{0},-\mathbf{1}), \ B^{\uparrow}=\begin{pmatrix}\mathbf{0}\\\mathbf{1}\end{pmatrix}, \ B^{\downarrow}=\begin{pmatrix}\mathbf{1}\\\mathbf{0}\end{pmatrix},$$

Matrix Product States

Matrix Product States: Coefficient of the wave function is the product of matrices. (gapped n-particle system)

Vertex: matrix internal line: inner product external line: spin indices

MERA

Tensor network of ground state wave-function of n-particle system in **quantum critical phase** can be written by **MERA** (Multi-scale entanglement renormalization ansatz) (G.Vidal 2007)





MERA and RG transformation



 τ direction: RG transformation (coarse-graining)



*** "disentangler**" which removes the short-range entanglement between each blocks (4-rank tensor)

Entanglement Entropy

Tensor Network is useful for the calculation of the entanglement entropy

Entanglement Entropy S_{EE} : Number of correlations between region A and region B.

 $S_{EE} = -Tr(\rho_A ln \rho_A)$

 $\rho_A = Tr_B \rho_{tot}$: reduced density matrix

 $\rho_{tot} = |\psi\rangle\langle\psi|$:density matrix of the total system



Entanglement Entropy of gapped system

Entanglement Entropy in (1+1)-dimensional gapped system is known to be constant (independent of system size *L*).



 $S_{EE} \sim const$

Entanglement Entropy by Matrix Product States



each matrix dimension: m

degree of freedoms at the two boundaries $= m \times m$

$$S_{EE} \sim ln m^2 \sim const$$

Entanglement Entropy in critical system

Entanglement entropy in (1+1)-dimensional critical system is not constant but has the **In L**-dependence for the long range correlation in the system.



 $S_{EE} \sim \ln L$

Entanglement entropy in critical phase



tensor dimension : *m*

Number of boundary bonds (blue bonds) $\sim ln L$

Total degree of freedoms at the boundary $\sim m^{lnL}$

Entanglement Entropy: $S_{EE} \sim \ln m^{lnL} \sim \ln L$

Entanglement Entropy by AdS/CFT

Entanglement Entropy by AdS/CFT (S. Ryu and T. Takayanagi '06)

$$S_{EE} = \frac{\gamma}{4G} \sim \ln L$$

- γ : the minimal length in **AdS3** space
- G: Newton constant



AdS space-time and MERA

MERA network \approx a discrete version of AdS space

(B. Swingle, 2009)

$$z=2^{\tau}$$







MERA network \approx a discrete version of AdS space

(B. Swingle, 2009)

MERA with periodic boundary condition

Anti de Sitter (AdS) space





by M. C. Escher

MERA at thermal system and AdS black hole

(d+1)-dimensional Quantum field theory

at thermal system







Thermo Field Dynamics Formalism

(Y. Takahashi and H. Umezawa, 1975)

the thermal state for temperature $\frac{1}{\beta}$

= the products of state in Hilbert space and that of copy (tilde) space

$$|\psi(\beta)\rangle = \frac{1}{\sqrt{Z(\beta)}} = e^{-\frac{\beta H}{2}}|I\rangle \quad |I\rangle = \sum_{n} |n\rangle|\tilde{n}\rangle = \sum_{n} |n\tilde{n}\rangle$$
$$H \dots \{|n\rangle\}$$

the Vacuum Expectation Value of the operator A

$$\langle A \rangle_s = \langle \psi(\beta) | A | \psi(\beta) \rangle = TrA\rho(\beta)$$

Tensor Network of Thermal State in single site model

Thermal state at $T = \frac{1}{2k_B\theta}$ can be written as an Matrix Product States.

$$|\boldsymbol{0}(\boldsymbol{\theta})\rangle = \left(\cos(\hbar\omega\boldsymbol{\theta}) + \sin(\hbar\omega\boldsymbol{\theta})\,a^{\dagger}\widetilde{a}^{\dagger}\,\right) \left|\boldsymbol{0}\widetilde{0}\rangle = \sum_{m,\widetilde{n}=0,1} A^{n}_{\alpha}A^{\widetilde{n}}_{\alpha}|n\widetilde{n}\rangle$$



$$A^{0} = \left(\sqrt{\cos(\hbar\omega\theta)}, 0\right) \quad A^{1} = \left(0, \sqrt{\sin(\hbar\omega\theta)}\right)$$
$$A^{\widetilde{0}} = \left(\frac{\sqrt{\cos(\hbar\omega\theta)}}{0}\right) \quad A^{\widetilde{1}} = \left(\frac{0}{\sqrt{\sin(\hbar\omega\theta)^{*}}}\right)$$

MERA at thermal system

By using Thermo Field Dynamics (Y. Takahashi and H. Umezawa, 1975), we suggest the following MERA at thermal system and the interface (red line) corresponds to the AdS black hole horizon.

$$|\psi\rangle = \sum_{\{n_j\}} \sum_{\{\widetilde{n}_k\}} A^{n_1 n_2 \cdots n_L} A^{\widetilde{n}_1 \widetilde{n}_2 \cdots \widetilde{n}_L} |n_1 n_2 \cdots n_L \widetilde{n}_1 \widetilde{n}_2 \cdots \widetilde{n}_L\rangle$$



Entanglement Entropy by thermal MERA

the degree of freedoms χ at the interface (red line)

 $\chi = m^A$

A: number of bonds at the interface

m: tensor dimensions



the entanglement entropy S_{EE} at the interface

L: system size

$$\frac{L}{2^{\tau_H}} = A \qquad S_{EE} = \ln \chi = \frac{L}{Z_H} \ln m$$

 $z=2^{\tau}$

 $z_H = 2^{\tau_H}$

Hawking Temperature

Entanglement Entropy in (1+1)-dimensional CFT at finite temperature,

$$S_{EE} = \frac{c}{3} \ln(\frac{\beta}{\pi\epsilon} \sinh\left(\frac{\pi L}{\beta}\right)) \simeq \frac{c}{3} \ln(\frac{\beta}{2\pi\epsilon}) + \frac{c}{3} \frac{\pi L}{\beta}$$

€: UV-cutoff (P. Calabrese and J. Cardy and J.S. Mech, 2004)

By comparing these two formula, we can find that

$$k_B T = \left(\frac{3}{c\pi} \ln m\right) \frac{1}{z_H}$$

Estimation of the tensor dimension

Hawking Temperature in AdS-black hole

$$k_B T = \frac{1}{2\pi z_H}$$

$$ds_{BH}^{2} = \frac{L^{2}}{z^{2}} \left(-f(z)dt^{2} + \frac{dz^{2}}{f(z)} + dx^{2} \right) \qquad f(z) = 1 - \left(\frac{z}{z_{H}}\right)^{2}$$

By comparing the Hawking temperature with the temperature given by MERA, we can get the relation between **central charge** and **tensor dimension** near the interface

$$\frac{3}{c\pi}\ln m = \frac{1}{2\pi} \quad \Longleftrightarrow \qquad m = e^{\frac{c}{6}}$$

Summary

Recently, the relation between AdS/CFT and MERA becomes interesting topic.

We consider that how the black hole horizon appears in the MERA network by using thermo filed dynamics (TFD) formalism and it is appeared as the interface between MERA and tilde-MERA network.

We can get the Hawking temperature scaling and the relation between central charge and tensor dimension.

Future Works

Relation between two-dimensional MERA and AdS/CFT (triangular lattice, square lattice...)

Relation between D-brane and top tensor of MERA (M. Nozaki, S.Ryu and T. Takayanagi, 2012).

